

Effects of multidisciplinary integrated care on quality of care in residential care facilities for elderly people: a cluster randomized trial

Marijke Boorsma MD, Dinnus H.M. Frijters PhD, Dirk L. Knol PhD, Miel E. Ribbe MD, Giel Nijpels MD, Hein P.J. van Hout PhD

See related commentary by Stadnyk and colleagues at www.cmaj.ca/cgi/doi/10.1503/cmaj.110789

ABSTRACT

Background: Sophisticated approaches are needed to improve the quality of care for elderly people living in residential care facilities. We determined the effects of multidisciplinary integrated care on the quality of care and quality of life for elderly people in residential care facilities.

Methods: We performed a cluster randomized controlled trial involving 10 residential care facilities in the Netherlands that included 340 participating residents with physical or cognitive disabilities. Five of the facilities applied multidisciplinary integrated care, and five provided usual care. The intervention, inspired by the disease management model, consisted of a geriatric assessment of functional health every three months. The assessment included use of the Long-term Care Facility version of the Resident Assessment Instrument by trained nurse-assistants to guide the design of an individualized care plan; discussion of outcomes and care priorities with the family physician, the resi-

dent and his or her family; and monthly multidisciplinary meetings with the nurse-assistant, family physician, psychologist and geriatrician to discuss residents with complex needs. The primary outcome was the sum score of 32 risk-adjusted quality-of-care indicators.

Results: Compared with the facilities that provided usual care, the intervention facilities had a significantly higher sum score of the 32 quality-of-care indicators (mean difference -6.7 , $p = 0.009$; a medium effect size of 0.72). They also had significantly higher scores for 11 of the 32 indicators of good care in the areas of communication, delirium, behaviour, continence, pain and use of antipsychotic agents.

Interpretation: Multidisciplinary integrated care resulted in improved quality of care for elderly people in residential care facilities compared with usual care.

Trial registration: www.controlled-trials.com trial register no. ISRCTN11076857.

Competing interests: None declared.

This article has been peer reviewed.

Correspondence to: Hein P.J. van Hout, hpj.vanhout@vumc.nl

CMAJ 2011; DOI:10.1503/cmaj.101498

The quality of care provided in residential care facilities is under pressure worldwide.¹ Facilities are frequently understaffed, and the complexity of care needed by residents increases while expertise of staff does not necessarily keep pace.^{2,3} Although most care organizations want to innovate and improve quality of care, many lack expertise or financial resources needed to do so.^{4,5} Family physicians are responsible for medical care in residential care facilities in the Netherlands. However, they do not regard themselves as suited for systematic management of chronic diseases and disabilities associated with frail health.⁶

About 10% of elderly people aged 75 or older in the Netherlands live in residential care facilities.^{7,8} These facilities were established to offer

sheltered living for elderly people who are disabled but still relatively healthy. Because of the growing elderly population, the characteristics of elderly people living in residential care facilities have become more comparable to those of people in nursing homes, who need complex care. Residential care facilities in the Netherlands are comparable to residential care facilities in Canada, are publicly funded and are subject to government inspection and approval. Over 70% of the residents need professional care, such as assistance with activities of daily living, nursing care (e.g., medication, wound care) and house-keeping. They have multiple chronic diseases and associated disabilities.⁹⁻¹²

Effective interventions for chronic illnesses generally rely on a multidisciplinary team ap-

proach. The elements of this approach include structured geriatric assessment, protocol-based regulation of medications, support for self-reliance and intensive follow-up. The closely related disease management model comprises coordination of care, steering of the care process and patient empowerment.¹³ This model is strongly recommended by Bodenheimer and colleagues to improve the health and quality of life of chronically ill patients.¹⁴ However, no studies have as yet been undertaken to evaluate the effects of disease management on functional health and quality of care for elderly people in residential care facilities who have physical or cognitive disabilities.

We developed an approach to multidisciplinary integrated care inspired by the disease management model. The objective of our study was to determine the effects of multidisciplinary integrated care on quality of care and quality of life for elderly people in residential care facilities.

Methods

The study was approved by the ethics committee of the VU University Medical Center in Amsterdam, the Netherlands. All participating residents or their proxies provided written informed consent. Figure 1 shows the flow of participants through the study.




Figure 1: Flow of completed interviews of participants through the trial. Because of missing data, the numbers of residents included in the primary and secondary analyses differed from the numbers interviewed. R = randomization.

Study design and participants

After a pilot study in one residential care facility, a cluster randomized controlled clinical trial was set up involving 10 facilities belonging to the umbrella care organization Omring, a large home care and long-term care provider in the Netherlands. The pilot facility was excluded from the trial. Randomization was carried out at the facility level; five facilities were assigned to the intervention group and the other five to the usual care group.

A total of 462 residents from the 10 facilities were recruited from December 2006 until December 2007. The average number of residents in each facility was 46, and staff included nurse-assistants and a house manager. All residents were listed in a primary care practice whose physician was responsible for their medical care. Residents who were terminally ill (as determined by staff or family physician) were excluded from the study.

Participating residents in each facility were visited by trained, blinded interviewers at baseline and at six months. If the resident was unable to understand the questions, a close family member was identified by staff and asked to act as proxy. The interview consisted of a computerized assessment of functional health, activities of daily living, depression, cognition, satisfaction with care, and use of medications. Proxies completed the interview except for the cognitive assessment, which was replaced by a short form of the Informant Questionnaire on Cognitive Decline in the Elderly.¹⁵

The duration of the trial was relatively short because of a high risk for dropout owing to the extreme vulnerability of residents and because the umbrella care organization intended to implement the care model in the control facilities as well. A detailed description of the design was reported earlier.¹⁶

Randomization

Randomization was carried out on facilities after matching for percentage of cognitively impaired residents, based on the assumption that a high percentage of such residents would affect care-related needs and services. In the matching procedure, the two facilities with the highest percentage of cognitively impaired residents were matched, and so on. Randomization was carried out using the first column from Pocock's random numbers table.¹⁷

Intervention

By adapting the principles of disease management, we introduced the concept of multidisciplinary integrated care. This concept focused on identification and monitoring of the functional

disabilities caused by chronic diseases. Its three basic elements correspond to those of the disease management model: monitoring of disabilities, coordination of care and empowerment.¹³ The third element is normally applied to patients only. However, we wanted to empower nurse-assistants in relation to monitoring and coordination of care, given that they provided all basic nursing care.

The model of multidisciplinary integrated care used in our study comprised five elements.¹⁶ First, a geriatric multidimensional assessment of all residents was conducted every three months. The Web-based Long-term Care Facility version 9.0 of the Resident Assessment Instrument was used for this purpose.¹⁸ The identified problem areas guide the design of an individualized care plan that is intended to improve or maintain functional health status. Second, the care plan was discussed with the resident, the resident's family and the family physician, and adapted to personal wishes. Third, residents with complex care needs were scheduled at least twice a year for a multidisciplinary meeting. Fourth, consultation with a geriatrician or psychologist was optional for the frailest residents with complex health care problems. Fifth, data from the Web-based Resident Assessment Instrument was used to provide an overview every three months of 32 risk-adjusted quality-of-care indicators. These indicators were compared with benchmark values derived from data on all residents of residential facilities in the Netherlands obtained using this instrument.^{19,20} Further details of the model of multidisciplinary integrated care are found in Appendix 1 (available at www.cmaj.ca/cgi/content/full/cmaj.101498/DC1).

Usual care

For facilities assigned to usual care, the family physician was responsible for medical care and offered it on request. There was neither coordination nor structured planning of care. Multidisciplinary meetings were mostly not attended by the family physicians.

Outcome measures

For the purpose of the evaluation, experienced, specially trained, blinded and supervised interviewers independently assessed the residents at baseline and six months later. The interviewers' assessments were supplemented by systematic observations by staff and extraction of data from residents' medical records (e.g., actual medication regimen).

Primary outcome measures

The first primary outcome was the sum score of the 32 risk-adjusted quality-of-care indicators,

which were developed by Morris²¹ and showed good validity and reliability. Appendix 2 (available at www.cmaj.ca/cgi/content/full/cmaj.101498/DC1) shows the definitions of the quality-of-care indicators, including their numerator and denominators. The quality-of-care indicators were originally based on observations recorded in the Long-term Care Facility assessment form. We incorporated the itemized observations needed to calculate these indicators in the assessments performed by our independently trained interviewers. Interrater reliability of the quality-of-care indicators between interviewers and nurse-assistants in the intervention facilities was satisfactory (mean intraclass correlation single measure 0.74). The sum score of the quality-of-care indicators was determined by the number of indicators per resident divided by the number of indicators applied to a resident. Indicators were calculated using mixed linear hierarchical models.

Four of the 32 quality-of-care indicators (behaviour problem, bladder or bowel incontinence, pressure ulcer and use of antipsychotics) applied both to the group of residents as a whole and to high- and low-risk groups. We therefore calculated two sum scores, one for all 32 indicators and one for the 24 whole-group indicators after exclusion of the 8 that were broken down to apply to high- and low-risk groups. Of these 24 whole-group indicators, on average 21 (standard deviation [SD] 6.6) applied to the residents. Of all 32 indicators, on average 22 (SD 6.1) applied to the residents. The relatively lower number of applicable indicators among all 32 indicators is due to the inclusion of the breakdown indicators, which applied to a maximum of 50% of the residents. The Cronbach's α of the sum score of the 24 whole-group indicators was 0.62.

For the second primary outcome, health-related quality of life was measured using a short-form 12-item version of the Rand Health Insurance Study questionnaire.²² Its properties were satisfactory when used by proxies, which was important in our study because of the percentage of cognitively impaired residents (58.2%).²³ We also calculated the number of quality-adjusted life-years using the algorithm of Hatoum and colleagues.²⁴

Secondary outcome measures

The secondary outcome measures comprised the 32 individual risk-adjusted quality-of-care indicators (described in Appendix 2); activities of daily living, as measured by the Groningen Activity Restriction Scale designed for elderly respondents and validated by Kempen and coauthors;²⁵ quality of care from the residents' per-

spective, as measured by a short (16-item) version of the QUOTE-Elderly instrument (Quality of Care from the Perspective of the Elderly);²⁶ hospital admissions recorded at the (single) local hospital, which covered more than 95% of all admissions;¹⁶ and mortality, as recorded by the interviewers or staff and cross-checked by the administration of the municipality.

Process outcomes

Process outcomes comprised the percentage of residents with completed assessments; the number of multidisciplinary meetings held, based on minutes of the meetings; the numbers of agreed-on medical, nursing and social actions, based on content analysis of care plans; and opinions of participating professionals regarding the intervention protocol, as obtained by interviews with staff and family physicians.

Statistical analysis

Sample-size calculations

Sample-size calculations were based on the expected effects of the intervention on quality of care and functional health using an α level of 0.05, a power of 80%, a dropout rate of 15% and an anticipated intracluster correlation of 0.05, based on Adams and colleagues,²⁷ across the residential facilities. To detect a fair benefit (i.e., Cohen's d effect size of 0.5), we estimated that the sample should include at least 82 residents in each of the two study groups.²⁸

Effect analyses

Effect analyses were performed according to both intention-to-treat and per-protocol principles. We accepted that the protocol was adhered to when the first two (obligatory) steps of the intervention were performed. We compared differences in the outcome measures over six months between the intervention and control groups using multivariable techniques. We applied mixed models for the continuous outcomes, using an unstructured covariance matrix for the longitudinal data. For the dichotomous outcomes, we applied generalized estimating equations using a logit link and an exchangeable working correlation.

In all effect analyses, we adjusted for baseline imbalance (e.g., by age, sex, cognitive impairment, depression, disability and interview by proxy). The analyses were done with multilevel specification if these variables showed significantly higher log-likelihood estimates. Because of our cluster randomized design (with randomization at the facility level), results of multilevel and "simple" analyses were compared for all outcomes. If higher-order clustering effects were

present, outcomes of the multilevel analyses were presented; if clustering was negligible, results of “simple” analyses were presented.

The intracluster coefficient across facilities was estimated by exchangeable working correlation for all individual (dichotomous) quality-of-care indicators. In all outcomes with pre-post measurements, the effect of interest was the group \times time (pre-post) interaction effect. A p value of 0.05 was considered to be significant.

Process analyses

We evaluated the extent to which the intervention program was performed according to protocol, the nature of the recommendations of the multidisciplinary meeting and the judgments of family physicians and staff about the intervention program.

Results

Sample and facility characteristics

Baseline characteristics of the residents and facilities are described in Table 1. There was a small imbalance between the intervention group and the usual care group in the numbers of residents with cognitive impairment and in the numbers with clinical depression. The trial ended up imbalanced because one control facility did not accept new entries as a result of an upcoming move to another location, and because a second control facility was in the middle of moving to a new building and could therefore recruit few residents for the study. Analyses without these facilities did not change the results.

Primary outcomes

Compared with residential care facilities that provided usual care, the intervention facilities had a significantly higher sum score of the 32 risk-adjusted quality-of-care indicators (mean difference -6.7 (95% confidence interval [CI] -8.69 to -4.71 , $p = 0.009$; Cohen d effect size 0.72) (Table 2). Self-reported quality of life did not differ between residents of control and intervention facilities (Table 3).

Secondary outcomes

The intervention facilities had higher scores than the control facilities for 30 of the 32 risk-adjusted indicators of quality of care; the scores for 11 of these 30 indicators had increased significantly (Table 2). In the intention-to-treat analyses, no differences in disability or quality of care as seen through residents' eyes were found between the two groups of facilities (Table 3). In the per-protocol analysis, residents in the intervention facilities tended to be more positive

about the quality of care over time than residents in the usual care facilities (difference 1.8 , $p = 0.072$). The per-protocol analyses showed that fewer residents died in the intervention group than in the control group (intervention $10/112$, control $25/139$; odds ratio 2.15 , 95% CI 1.06 – 4.38 ; $p = 0.035$).

Table 1: Baseline characteristics of the 10 residential care facilities and the 340 participating residents

Characteristic	Intervention group	Control group
Facilities	<i>N</i> = 5	<i>N</i> = 5
No. of participating residents, mean (SD)	39.8 (17.7)	27.2 (18.0)
No. of personnel, full-time equivalent per resident, mean (SD)	0.33 (0.04)	0.44 (0.10)
Residents	<i>n</i> = 201	<i>n</i> = 139
Age, yr, mean (SD)	85.8 (6.2)	85.5 (8.0)
Sex, male, no. (%)	48 (23.9)	36 (25.9)
Marital status, no. (%)	<i>n</i> = 191	<i>n</i> = 135
Married	42 (22.0)	27 (20.0)
Widowed	130 (68.1)	93 (68.9)
Never married	19 (9.9)	15 (11.1)
Education, no. (%)	<i>n</i> = 190	<i>n</i> = 132
Primary school or less	112 (58.9)	79 (59.8)
Lower technical vocational training	45 (23.7)	26 (19.7)
Average and higher vocational training	34 (17.9)	30 (22.7)
Clinical characteristics, no. (%)	<i>n</i> = 199	<i>n</i> = 136
Asthma or COPD	29 (14.6)	13 (9.6)
Myocardial infarction	61 (30.7)	30 (22.1)
Cerebrovascular accident	33 (16.6)	29 (21.3)
Diabetes	39 (19.6)	27 (19.9)
Arthritis	73 (36.7)	47 (34.6)
Cancer	12 (6.0)	11 (8.1)
Hypertension	35 (17.6)	32 (23.5)
Chronic somatic diseases, no., mean (SD)	1.54 (1.25)	1.49 (1.17)
Cognitive impairment*	101 (50.8)	90 (66.2)
Clinical depression†	10 (5.0)	16 (11.8)
Body mass index, mean (SD)	<i>n</i> = 164 26.2 (4.98)	<i>n</i> = 116 26.3 (4.67)
Falls experienced in past 30 days, no. (%)	<i>n</i> = 190	<i>n</i> = 134
One	23 (12.1)	18 (13.4)
Two or more	8 (4.2)	10 (7.5)
Total	31 (16.3)	28 (20.9)
Medications, no., mean (SD)	7.6 (3.4)	8.0 (3.6)
4-m walking time, sec, median (IQR)	4.0 (1.0–7.75)	5.0 (1.0–8.0)
Not able to do walking test	<i>n</i> = 172 76 (44.2)	<i>n</i> = 114 59 (51.8)

Note: COPD = chronic obstructive pulmonary disease, IQR = interquartile range, SD = standard deviation.

*Score of < 5 on the Memory Impairment Screen, or ≥ 3.6 on the short form of the Informant Questionnaire on Cognitive Deterioration.

†Diagnosed by family physician or specialist.

Table 2: Risk-adjusted indicators of quality of care for elderly people in intervention and control residential care facilities during the six-month study period

Indicator	Intraclass coefficient	Group; % of residents		Adjusted OR* (95% CI)
		Intervention facilities	Control facilities	
Worsening of activities of daily living	0.02	14.7	48.5	0.17 (0.07–0.40)
Worsening of locomotion	< 0.001	16.1	7.1	2.85 (0.91–8.96)
Increase in no. of falls	–0.03	17.6	9.6	2.00 (0.69–5.64)
Decline in cognitive ability	0.03	18.4	24.1	0.84 (0.51–1.38)
Decline in communication	0.01	17.5	46.9	0.25 (0.13–0.49)
Delirium, new or persistent	0.06	28.2	56.3	0.27 (0.10–0.69)
Behavioural problem	0.01	8.7	26.5	0.27 (0.10–0.74)
High risk	0.04	15.4	45.5	0.24 (0.07–0.89)
Low risk	0.04	2.0	8.6	0.20 (0.03–1.34)
Little or no social activity	–	0.0	0.0	–
New in-dwelling catheter	–0.03	1.0	4.6	0.04 (0.006–0.29)
In-dwelling catheter	0.01	2.0	0.0	–
Bladder or bowel incontinence	–0.04	33.3	46.2	0.78 (0.66–0.91)
High risk	–0.06	62.5	71.1	0.79 (0.28–2.28)
Low risk	–0.06	19.4	34.1	0.52 (0.40–0.67)
Decline in bowel continence	–0.03	6.9	23.4	0.25 (0.15–0.45)
Decline in bladder continence	–0.02	14.6	45.9	0.17 (0.07–0.38)
Urinary tract infection	0.01	1.9	4.5	0.79 (0.25–2.49)
Use of feeding tube	< 0.001	0.0	1.5	–
Low body mass index	< 0.001	1.9	0.0	–
Weight loss	0.001	3.9	4.6	0.67 (0.09–5.00)
Inadequate pain management	–0.01	21.4	13.2	1.81 (0.91–3.57)
Worsening of pain	–0.03	12.9	40.9	0.20 (0.07–0.52)
Presence of pressure ulcers	–0.01	4.9	7.5	0.63 (0.21–1.91)
High risk	–0.17	16.0	19.0	0.80 (0.18–3.44)
Low risk	0.02	1.3	2.1	0.51 (0.06–4.04)
Worsening of pressure ulcers	0.03	1.9	5.2	0.57 (0.31–1.05)
Burns, skin tears or cuts	0.02	8.7	4.5	1.9 (0.91–4.14)
Use of physical restraints	–	0.0	1.5	–
Use of antipsychotic agents	0.01	3.8	11.0	0.25 (0.08–0.78)
High risk	17.00	25.0	25.0	0.89 (0.03–36.1)
Low risk	0.01	2.1	10.7	0.15 (0.03–0.66)
Sum score				Mean difference (95% CI)
For all 32 quality-of-care indicators, range 0%–100% (SD)†‡	0.01	11.5 (9.0)	18.2 (9.4)	–6.7 (–8.69 to –4.71)¶
For 24 whole-group quality-of-care indicators, range 0%–100% (SD)†§	0.05	9.1 (9.4)	15.7 (11.2)	–6.6 (–8.81 to –4.39)**

Note: CI = confidence interval, OR = odds ratio, SD = standard deviation.
 *Calculated by generalized linear models. ORs and sum scores for quality-of-care risk indicators were adjusted for sex, age, interview by proxy, depression and cognitive impairment at baseline, accounting for within-facility clustering.
 †The sum score was determined by the number of quality-of-care indicators per resident divided by the number of quality-of-care indicators applied to a resident. Indicators were calculated using mixed linear hierarchical models.
 ‡Of the 32 quality-of-care indicators, a mean of 23.1 applied to residents in intervention facilities and 20.1 applied to residents in control facilities.
 §The 24 whole-group quality-of-care indicators excludes the 8 indicators broken down to apply to high- and low-risk groups. Of the 24 whole-group indicators, a mean of 21.7 applied to residents in intervention facilities and 19.5 applied to residents in control facilities.
 ¶ $p = 0.009$.
 ** $p = 0.021$.

Process of care

The first step of the protocol — assessment with the Long-term Care Facility version of the Resident Assessment Instrument — was completed for 55.2% of the residents in the intervention facilities. This proportion was less than we had aimed for and was partly a result of implementation delay.

Forty multidisciplinary meetings were held in the intervention facilities during the study period, and the outcomes of assessment of 93 residents included in the study were discussed (Table 4). The primary care physician was present at 90% of the multidisciplinary meetings, and the geriatrician at 75%. By contrast, only 25% of the multidisciplinary meetings in the control facilities were visited by the primary care physician. The number of recommended actions per resident was 3.67 in the intervention facility meetings and 2.26 in the control facility meetings.

The expertise of staff was felt by 52.9% of staff and 54.5% of the family physicians to have increased after introduction of the care model. The same percentage of staff and 63.6% of family physicians felt that they had more knowledge about residents' health. In addition, 58.8% of staff and 81.8% of the family physicians felt that their cooperation had improved. About 55% of the family physicians considered quality of care to have improved; 73% acknowledged the need for a new care model.

Ancillary analyses

We did not find effect modification of the outcomes by age, sex or baseline disability.

Interpretation

Compared with usual care, our model of multidisciplinary integrated care resulted in substantially higher quality of care for elderly people in residential care facilities. Functional ability, number of hospital admissions and health-related quality of life remained comparable between the two groups. According to the per-protocol analyses, mortality was lower in the intervention facilities and residents in the intervention facilities were more positive about their quality of care. Owing to the short intervention period (six months), the full protocol was applied to less than half of the residents in the intervention facilities. The training and empowerment of nurse-assistants, which was completed for all intervention facilities, together with monitoring using the geriatric assessment instrument, were likely to be the most important ingredients for improvement of the quality of care.

Earlier studies have reflected elements of our intervention. For example, positive health effects on residents have been reported as a result of interdisciplinary geriatric primary care in American facilities.²⁹ Integrated and home-based geriatric care management was reported to improve quality of care and reduce use of acute care ser-

Table 3: Health-related outcomes and residents' opinions of quality of care

Outcome measure	Intracluster coefficient	Intervention group	Control group	Effect statistic	p value
Continuous (range)		Mean (SD)		Group × time	
Short 12-item version of the Rand Health Insurance Study questionnaire (0–100)	0.02			1.02	0.35
Baseline		43.34 (5.96)	42.33 (6.86)		
6 mo		42.31 (6.04)	42.56 (6.35)		
Quality-adjusted life-years (0–1)	0.05				
6 mo		0.28 (0.11)	0.27 (0.12)	0.87	
Groningen Activity Restriction scale (18–72)	0.13			–1.40	0.16
Baseline		43.70 (13.73)	39.90 (13.91)		
6 mo		42.41 (13.37)	39.06 (13.70)		
Quality of care through residents' eyes (16–64)	0.07			1.56	0.12
Baseline		57.41 (5.46)	58.01 (6.69)		
6 mo		56.32 (6.47)	56.10 (6.64)		
Dichotomous		No. (%)		OR (95% CI)	
Mortality	–0.02	28/201 (13.9)	25/139 (18.0)	1.09 (0.87–1.38)	0.44
Admissions to hospital ≥ 1	–0.02	22/142 (15.5)	12/85 (14.1)	1.32 (0.94–1.87)	0.11

Note: CI = confidence interval, OR = odds ratio, SD = standard deviation.

Table 4: Characteristics and outcomes of multidisciplinary meetings held during the six-month study period

Characteristic or outcome	Intervention facility meetings n = 40	Control facility meetings n = 28
Meetings at which primary care physician was present, no. (%)	36 (90)	7 (25)
Meetings at which geriatrician was present, no. (%)	30 (75)	21 (75)
Meetings at which psychologist was present, no. (%)	21 (53)	16 (57)
No. of residents discussed, total (per meeting)	93 (2.1)	68 (2.4)
No. of actions on care plan, total (per resident)		
Medical	92 (0.99)	60 (0.88)
Nurse care	124 (1.33)	27 (0.40)
Referral to medical specialist	32 (0.34)	12 (0.18)
Paramedical referral	53 (0.57)	37 (0.41)
Medication change	40 (0.43)	18 (0.25)
All	341 (3.67)	154 (2.26)

vices in a high-risk group of low-income elderly people living at home.^{30,31} Use of the Home Care version of the Resident Assessment Instrument in primary care health centres in Hong Kong resulted in improvement in 2 of 13 functional outcomes.³² Modest positive effects on well-being and on deterioration of functional skills were found in a multidisciplinary program in vulnerable older people living at home.³³

Limitations

Our study was limited by the fact that our population was frail and comprised a high percentage of cognitively impaired residents. As a result, a portion of the data was collected from interviews with proxies. The judgments of proxies may have differed from the residents' judgments. Therefore, we adjusted for proxy interview and cognitive status in our analyses. The cluster randomization produced an imbalance between the intervention and control facilities in the number of participating residents and in some of the functional characteristics of the residents at baseline. Although we adjusted for the imbalance in functional characteristics, imbalance in the number of participating residents may have led to underpowered results.

Variation across the intervention facilities in the application of the complete protocol (3%–66%) was another limitation. This variation can be explained by financial and administrative issues during the study period. The financial obligations for residential care facilities resulting from a new

national funding system for residential care of elderly people caused uncertainty about job continuation, high turnover of managers and new priorities at the facilities in our study. Despite this limitation, the improvement in quality of care at the facilities in our study was substantial.

Conclusion

Our model of multidisciplinary integrated care resulted in improved quality of care for elderly people in residential care facilities compared with usual care. The results of this study are applicable to elderly people in such settings as residential care facilities and nursing homes, and even elderly people living in the community. In primary care settings, it may be beneficial to have a model to monitor elderly people and those with chronic diseases, to prevent functional decline and admission to hospital for acute care. It is also important to have an instrument that not only delivers output on the patient level but also on the management level, to facilitate monitoring of quality of care by managers in a sector of health care that is under enormous societal pressure to improve its performance.

References

1. Bos JT, Frijters DH, Wagner C, et al. Variations in quality of home care between sites across Europe, as measured by home care quality indicators. *Aging Clin Exp Res* 2007;19:323-9.
2. Broese van Groenou MI. Unequal chances for reaching "a good old age." Socio-economic health differences among older adults from a life course perspective [article in Dutch]. *Tijdschr Gerontol Geriatr* 2003;34:196-207.
3. Spector WD, Cohen JW, Pesis-Katz I. Home care before and after the Balanced Budget Act of 1997: shifts in financing and services. *Gerontologist* 2004;44:39-47.
4. Challis D, Hughes J. Residential and nursing home care — issues of balance and quality of care. *Int J Geriatr Psychiatry* 2003;18:201-4.
5. Challis D, Stewart K, Donnelly M, et al. Care management for older people: Does integration make a difference? *J Interprof Care* 2006;20:335-48.
6. Challis D, Hughes J. Frail old people at the margins of care: some recent research findings. *Br J Psychiatry* 2002;180:126-30.
7. Pot AM, Portrait F, Visser G, et al. Utilization of acute and long-term care in the last year of life: comparison with survivors in a population-based study. *BMC Health Serv Res* 2009;9:139.
8. Tomassini C, Glaser K, Wolf DA, et al. Living arrangements among older people: an overview of trends in Europe and the USA. *Popul Trends* 2004;Spring:24-34.
9. Geerlings SW, Pot AM TJ, Deeg DJG. Predicting transitions in the use of informal and professional care by older adults. *Ageing Soc* 2005;25:111-30.
10. Simonsick EM, Kasper JD, Phillips CL. Physical disability and social interaction: factors associated with low social contact and home confinement in disabled older women (the Women's Health and Aging Study). *J Gerontol B Psychol Sci Soc Sci* 1998; 53:S209-17.
11. Sprangers MA, de Regt EB, Andries F, et al. Which chronic conditions are associated with better or poorer quality of life? *J Clin Epidemiol* 2000;53:895-907.
12. Strawbridge WJ, Sheme SJ, Balfour JL, et al. Antecedents of frailty over three decades in an older cohort. *J Gerontol B Psychol Sci Soc Sci* 1998;53:9-16.
13. Bijl D. Effectiveness of disease management programmes for recognition, diagnosis and treatment of depression in primary care. *Eur J Gen Pract* 2004;10:6-12.
14. Bodenheimer T, Wagner EH, Grumbach K. Improving primary care for patients with chronic illness: the chronic care model, Part 2. *JAMA* 2002;288:1909-14.
15. Jorm AF. A short form of the Informant Questionnaire on Cog-

- nitive Decline in the Elderly (IQCODE): development and cross-validation. *Psychol Med* 1994;24:145-53.
16. Boorsma M, van Hout HP, Frijters DH, et al. The cost-effectiveness of a new disease management model for frail elderly living in homes for the elderly: design of a cluster randomized controlled clinical trial. *BMC Health Serv Res* 2008;8:143.
 17. Pocock S. *Clinical trials: a practical approach*. New York (NY): Wiley and Sons; 1983.
 18. Hirdes JP, Fries BE, Morris JN, et al. Integrated health information systems based on the RAI/MDS series of instruments. *Healthc Manage Forum* 1999;12:30-40.
 19. Frijters D, Gerritsen D, Steverink N, et al. Care quality: reliability and usefulness of observation data in bench marking nursing homes and homes for the aged in the Netherlands. *Tijdschr Gerontol Geriatr* 2003;34:21-9.
 20. Zimmerman DR. Improving nursing home quality of care through outcomes data: the MDS quality indicators. *Int J Geriatr Psychiatry* 2003;18:250-7.
 21. Morris J. *Validation of long-term and post-acute care quality indicators, final report*. Cambridge (MA): Abt Associates Inc.; Rosalind (MA): HRCR Research and Training Institute; Providence (RI): Brown University Center for Gerontology and Health Care Research; 2003.
 22. Brook RH, Ware JE Jr, Davies-Avery A, et al. Overview of adult health status measures fielded in Rand's Health Insurance Study. *Med Care* 1979;17(Suppl 7):1-131.
 23. Brazier JE, Harper R, Jones NM, et al. Validating the SF-36 health survey questionnaire: new outcome measure for primary care. *BMJ* 1992;305:160-4.
 24. Hatoum HT, Brazier JE, Akhras KS. Comparison of the HUI3 with the SF-36 preference based SF-6D in a clinical trial setting. *Value Health* 2004;7:602-9.
 25. Kempen GI, Miedema I, Ormel J, et al. The assessment of disability with the Groningen Activity Restriction Scale. Conceptual framework and psychometric properties. *Soc Sci Med* 1996;43:1601-10.
 26. Sixma HJ. Quality of care from the perspective of elderly people: the QUOTE-Elderly Instrument. *Age Ageing* 2000;29:173-8.
 27. Adams G. Patterns of intra-cluster correlation from primary care research to inform study design and analysis. *J Clin Epidemiol* 2004;57:785-94.
 28. Cohen J. *Statistical power analysis for the behavioural sciences*. New York (NY): Academic Press; 1977.
 29. Burns R, Nichols LO, Graney MJ, et al. Impact of continued geriatric outpatient management on health outcomes of older veterans. *Arch Intern Med* 1995;155:1313-8.
 30. Counsell SR, Callahan CM, Clark DO, et al. Geriatric care management for low-income seniors: a randomized controlled trial. *JAMA* 2007;298:2623-33.
 31. Counsell SR, Callahan CM, Tu W, et al. Cost analysis of the Geriatric Resources for Assessment and Care of Elders care management intervention. *J Am Geriatr Soc* 2009;57:1420-6.
 32. Chi I, Chou KL, Kwan CW, et al. Use of the Minimum Data Set — Home Care: a cluster randomized controlled trial among the Chinese older adults. *Aging Ment Health* 2006;10:33-9.
 33. Melis RJ, van Eijken MI, Borm GF, et al. The design of the Dutch EASYcare study: a randomised controlled trial on the effectiveness of a problem-based community intervention model for frail elderly people [NCT00105378]. *BMC Health Serv Res* 2005;5:65.

Affiliations: From the Departments of General Practice (Boorsma, Nijpels, van Hout), Nursing Home Practice (Frijters, Ribbe), and Epidemiology and Biostatistics (Knol), EMGO Institute for Health and Care Research, VU University Medical Center, Amsterdam, the Netherlands

Contributors: Marijke Boorsma, Hein van Hout and Giel Nijpels were involved in the conception and design of the study. Marijke Boorsma and Dinnus Frijters were involved in the acquisition of the data. Marijke Boorsma, Hein van Hout, Dirk Knol and Miel Ribbe were involved in the analysis and interpretation of the data. Marijke Boorsma and Hein van Hout were involved in the drafting of the article. All of the authors were involved in the critical revision of the manuscript for important intellectual content and approved the final version submitted for publication.

Funding: This study was funded by the Netherlands Organization for Health Research and Development (ZonMw grant no. 945-05-030). The funding organization had no role in the design and conduct of the study; the collection, management, analysis and interpretation of the data, or the preparation, review and approval of the manuscript.